

TITLE:	Form, Autologous Donor Order		
DEPARTMENT:	03-Special Donations	REF #	03-01-05
FORM #:	03-F66	FORM/VER:	2
		PAGE:	1 of 1

Autologous Donor Order Form

Ref: 03-01-05

Appointments and Information: (650) 723-6667 / FAX: (650) 723-8155 / EMAIL: specialdonations@stanford.edu

Incomplete Orders Will Not Be Accepted

Number and Type of Units Requested:		FOR INTERNAL SD DEPT. USE ONLY:
Packed Red Blood Cells		CPD/AS1
Other:		

Patient Name: _____ Date of Birth: _____

Last First MI

Medical Record Number: _____ Phone: Day: _____ Eve: _____

Type of Procedure Scheduled: _____ Gender: Male ☐ Female ☐

☐ Surgery ☐ Transfusion Date: _____ ICD Code: _____

Location for Transfusion: ☐ SHC ☐ LPCH ☐ Other:

PHYSICIAN AUTHORIZATION FOR AUTOLOGOUS COLLECTION

Note: Stanford/LPCH Transfusion Service will NOT accept autologous units from patients known to be infected with HIV, HCV, or HBV, or positive on the blood bank testing for the following infectious agents: HIV, HBV, HCV, WNV, HTLV I/II, ZIKV and/or *T. cruzi*.

Blood donation involves an acute loss of 10-15% of the patient's blood volume and may cause transient hypotension. Stanford Blood Center (SBC) requires authorization by a physician familiar with the patient's condition prior to performing phlebotomy for patients with the following conditions. Note that the failure to provide authorization at the time of ordering will cause delays in donation.

SBC will make the final decision on donor eligibility at the time of donation. Patients with the following conditions will not be drawn:

- Angina at REST within the past three months
- Requirement for supplemental O₂ except those with secondary polycythemia

I am familiar with the patient's condition and authorize SBC to perform one or more blood draws for autologous blood collection. This patient has a history of the following (check all that apply):

- ☐ Current heart valve disease or congestive heart failure
- ☐ History of angina, MI, stroke, TIA, or heart surgery within the past year
- ☐ Pregnancy, currently or within the past six weeks
- ☐ Sickle Cell Trait (HqSA)

To minimize the risk of pre-operative anemia, SBC currently requires autologous donors to have a minimum hemoglobin level of 12.5 g/dL for women and 13.0 g/dL for men. However, as the ordering physician, you can consent to the lower acceptable limit of 11.0 g/dL by indicating below:

- The lower hemoglobin limit at the time of donation is 12.5 g/dL for women and 13.0 g/dL for men.
- The lower hemoglobin limit at the time of donation is 11.0 g/dL.

Print Physician Name: _____ Physician Signature: _____

Date: Physician Address:

Physician Phone: _____ Physician Fax: _____

FOR BLOOD CENTER USE ONLY

Comments:

Physician Contact Info. Verified by: _____ Date: _____